

AAP-Commissioned Review

Lasers in Periodontics: A Review of the Literature

Charles M. Cobb*

Background: Despite the large number of publications, there is still controversy among clinicians regarding the application of dental lasers to the treatment of chronic periodontitis. The purpose of this review is to analyze the peer-reviewed research literature to determine the state of the science concerning the application of lasers to common oral soft tissue problems, root surface detoxification, and the treatment of chronic periodontitis.

Methods: A comprehensive computer-based search combined the following databases into one search: Medline, Current Contents, and the Cumulated Index of Nursing and Allied Health. This search also used key words. In addition, hand searches were done for several journals not cataloged in the databases, and the reference lists from published articles were checked. All articles were considered individually to eliminate non-peer-reviewed articles, those dealing with commercial laser technology, and those considered by the author to be purely opinion articles, leaving 278 possible articles.

Results: There is a considerable conflict in results for both laboratory studies and clinical trials, even when using the same laser wavelength. A meaningful comparison between various clinical studies or between laser and conventional therapy is difficult at best and likely impossible at the present. Reasons for this dilemma are several, such as different laser wavelengths; wide variations in laser parameters; insufficient reporting of parameters that, in turn, does not allow calculation of energy density; differences in experimental design, lack of proper controls, and differences in severity of disease and treatment protocols; and measurement of different clinical endpoints.

Conclusions: Based on this review of the literature, there is a great need to develop an evidence-based approach to the use of lasers for the treatment of chronic periodontitis. Simply put, there is insufficient evidence to suggest that any specific wavelength of laser is superior to the traditional modalities of therapy. Current evidence does suggest that use of the Nd:YAG or Er:YAG wavelengths for treatment of chronic periodontitis may be equivalent to scaling and root planing (SRP) with respect to reduction in probing depth and subgingival bacterial populations. However, if gain in clinical attachment level is considered the gold standard for non-surgical periodontal therapy, then the evidence supporting laser-mediated periodontal treatment over traditional therapy is minimal at best. Lastly, there is limited evidence suggesting that lasers used in an adjunctive capacity to SRP may provide some additional benefit. *J Periodontol* 2006;77:545-564.

KEY WORDS

Bacteria; calculus; chronic periodontitis; lasers; periodontics; root.

Periodically, the Board of Trustees of the American Academy of Periodontology identifies the need for review of the literature on a specific topic and requests the Editor-in-Chief of the Journal of Periodontology to commission such a review. The selected author is solely responsible for the content, and the manuscript is peer reviewed, like all other Journal articles. The Academy's Board of Trustees does not review or approve the manuscript prior to publication, and the content of the review should not be construed as Academy policy.

Based on Albert Einstein's theory of spontaneous and stimulated emission of radiation, Maiman developed the first laser prototype in 1960.¹ Maiman's device used a crystal medium of ruby that emitted a coherent radiant light from the crystal when stimulated by energy. Thus, the ruby laser was created. Shortly thereafter, in 1961, Snitzer² published the prototype for the Nd:YAG laser. The first application of a laser to dental tissue was reported by Goldman et al.³ and Stern and Sognnaes,⁴ each article describing the effects of the ruby laser on enamel and dentin. However, the current relationship of

* Department of Periodontics, School of Dentistry, University of Missouri, Kansas City, MO.

dentistry with the laser takes its origins from an article published in 1985 by Myers and Myers⁵ describing the in vivo removal of dental caries using a modified ophthalmic Nd:YAG laser.⁴ Four years later, it was suggested that the Nd:YAG laser could be used for oral soft tissue surgery,⁶ which ultimately lead to the present relationship between lasers and clinical periodontics.⁷⁻⁹

The subject of lasers in periodontics now encompasses a rapidly increasing and significant volume of published literature. Despite the large number of publications, there is still controversy among clinicians regarding the application of dental lasers to the treatment of periodontal diseases, and more specifically, chronic periodontitis. The purpose of this review is to analyze the peer-reviewed research literature to determine the state of the science regarding the application of lasers to common oral soft tissue problems, root surface detoxification, and the treatment of chronic periodontitis.

MATERIALS AND METHODS

The search to locate relevant articles for this review was undertaken at several levels. The first two searches were done with Medline-Ovid using an explode of “periodontal diseases” and an explode of “lasers.” The result of this search was first limited to English only and randomized clinical trials, which resulted in 23 articles.

A comprehensive second search combined the following databases into one search: Medline, Current Contents, and the Cumulated Index of Nursing and Allied Health. This search combined the truncated search words: periodont\$ or gingiv\$, or mouth mucosa and laser\$. Duplicates were removed, leaving 1,137 possible articles. This result was limited to English-only articles and the years 1990 to 2005, for a total of 906 articles. In addition, hand searches were done for the *Journal of Oral Laser Applications*, the proceedings of the Lasers in Dentistry conferences sponsored by the International Society for Optical Engineering (SPIE), and the Proceedings of the International Congress on Lasers in Dentistry. Lastly, the reference lists from published articles were checked. All articles were considered individually to eliminate non-peer-reviewed articles, those dealing with commercial laser technology, and articles considered by the author to be purely opinion articles, leaving 278 possible articles.

In compliance with the *Journal of Periodontology* instructions to authors, free-standing abstracts were not considered for inclusion in this review. From this final number of articles, the majority of those cited in this review were published since 1995. It should be noted that a few selected articles published as literature reviews or opinion articles or those concerned with the physics of laser-tissue interactions were used

to present background material and to place comments by the present author in context.

Table 1 shows the results of the literature search by journal title. Table 2 displays the number of studies categorized by their experimental design. In order of decreasing clinical relevance, the experimental designs were as follows: randomized, blinded, controlled, longitudinal, clinical trials, cohort or longitudinal studies,

Table 1.

Results of the Computer-Based Literature Search by Journal Title for Articles on Lasers and Periodontics

Journal Name	Number of Published Articles (1990 to 2005)
<i>Compendium of Continuing Dental Education</i>	5
<i>Dental Clinics of North America</i>	13
<i>Dental Economics</i>	8
<i>Dentistry Today</i>	15
<i>International Journal of Periodontics & Restorative Dentistry</i>	3
<i>Journal of the American Dental Association</i>	3
<i>Journal of Clinical Laser Medicine and Surgery</i>	7
<i>Journal of Clinical Periodontology</i>	12
<i>Journal of Dentistry</i>	1
<i>Journal of Dental Research</i>	2
<i>Journal of the International Academy of Periodontology</i>	3
<i>Journal of Oral Laser Applications</i>	22
<i>Journal of Periodontology</i>	39
<i>Journal of Periodontics and Esthetic Dentistry</i>	1
<i>Journal of Periodontal Research</i>	5
<i>Lasers in Dentistry: Proceedings of SPIE</i>	25
<i>Lasers in Surgery and Medicine</i>	30
<i>Periodontology 2000</i>	2
<i>Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics</i>	2
<i>Quintessence International</i>	1
Miscellaneous	79
Total number of published articles	278

Table 2.
Number of Published Articles on Lasers in Periodontics Listed by Experimental Design in Order of Decreasing Clinical Relevance

Experimental Design	Number of Published Articles	Percentage of Total
Randomized, blinded, controlled, longitudinal, clinical trials	3	1.1
Cohort or longitudinal studies	20	7.2
Case-controlled studies	12	4.3
Non-controlled case studies	25	9.0
Descriptive studies	44	15.8
In vivo animal studies	27	9.7
In vitro laboratory studies	59	21.2
Reviews of the literature	88	31.7
Total number of articles	278	100.0

case-controlled studies, non-controlled case studies, descriptive studies, in vivo animal studies, and in vitro laboratory studies.¹⁰ The category of literature review has been added for informational purposes.

Importance of Wavelength

Typically, lasers are named according to the active element(s) that is induced to undergo the stimulated quantum transitions that, in turn, creates the energy beam. Thus, lasers commonly used in dentistry consist of a variety of wavelengths delivered as either a continuous, pulsed (gated), or running pulse waveform, e.g., CO₂, Nd:YAG, Ho:YAG, Er:YAG, Er,Cr:YSGG, Nd:YAP, GaAs (diode), and argon (Table 3).

The following description pertains to those laser wavelengths and waveforms currently available to dentistry. Shorter wavelengths and pulse widths combined with higher-power densities as seen with some medical and industrial lasers bring with them other interaction phenomena that are not currently relevant to dental applications and, therefore, will not be discussed in this article.

The energy emitted by a laser is essentially a light of one color (i.e., monochromatic) and, therefore, of one wavelength. The photons comprising the energy beam are emitted as a coherent (in phase), unidirectional, monochromatic light that can be collimated into an intensely focused beam that exhibits little divergence. The focused energy beam will interact with a target material by being absorbed, reflected, or scattered. In the case of biologic tissues, the laser energy is

absorbed by the target surface tissues and will only exhibit scattering in cases of deep tissue penetration. The absorbed light energy is converted to heat and constitutes a photothermal event. Depending on various parameters, the absorbed energy can result in simple warming, coagulation, or excision and incision through tissue vaporization. Variable parameters affecting energy absorption include emission wavelength, power (watts), waveform (continuous or pulsed), pulse duration, energy/pulse, energy density, duration of exposure, peak power of pulse, angulation of the energy delivery tip to the target surface, and optical properties of the tissue.

Although the wavelength of light is the primary variable that determines the extent of energy absorption by a target tissue, one must also be aware of the optical properties of the tissue. The optical properties of a tissue dictate, to a great extent, the interaction with specific laser wavelengths. For example, optical properties of tissues comprising the periodontium include such factors as pigmentation, water content, mineral content, heat capacity that accounts for both thermal conductivity and tissue density, and latent heats of transformation (i.e., denaturing of proteins, vaporization of water, and melting of mineral). Bone is considered the classic composite tissue, being comprised of $\approx 67\%$ inorganic minerals (calcium hydroxyapatite) and 33% collagen and non-collagenous proteins. By contrast, gingiva is comprised of varying densities of fibrous connective tissue, associated extracellular matrix components, and a high content of water ($\approx 70\%$). Additionally, the gingivae frequently exhibit melanin pigmentation. Other factors that likely play a role in laser-tissue interactions include the physiologic and mechanical processes of heat conduction and dissipation, the degree of tissue inflammation and vascularity, and the availability of progenitor cells to participate in the healing process.

Each wavelength of laser energy is absorbed to a greater or lesser degree in water, pigment, or hydroxyapatite. As examples, the CO₂ laser (10,600-nm wavelength) has a high absorption coefficient in water and consequently is well suited for soft tissue surgery but currently has no scientifically well-supported clinical application to mineralized tissues. The Nd:YAG (1,064-nm wavelength) and diode lasers (800- to 950-nm wavelength) have lower absorption coefficients in water than CO₂ lasers but are preferentially absorbed in pigmented tissues, and the Er,Cr:YSGG and Er:YAG wavelengths (2,780 and 2,940 nm, respectively) are highly absorbed in both water and hydroxyapatite. Given the diversity of available wavelengths, the prudent clinician should first determine the specific clinical treatment goals and then select the technology (laser or otherwise) best suited to achieve the desired endpoint(s).

Table 3.
Characteristics of Laser Wavelengths Used in Clinical Dentistry

Laser Type	Common Abbreviation	Wavelength	Waveform	Delivery Tip	Reported Periodontal Applications
Carbon dioxide	CO ₂	10.6 μm	Gated or continuous	Hollow waveguide; beam focused when 1 to 2 mm from target surface	Soft tissue incision and ablation; subgingival curettage
Neodymium:yttrium-aluminum-garnet	Nd:YAG	1.064 μm	Pulsed	Flexible fiber optic system of varying diameters; surface contact required for most procedures	Soft tissue incision and ablation; subgingival curettage and bacterial elimination
Holmium:yttrium-aluminum-garnet	Ho:YAG	2.1 μm	Pulsed	Flexible fiber optic system; surface contact required for most procedures	Soft tissue incision and ablation; subgingival curettage and bacterial elimination
Erbium:yttrium-aluminum-garnet	Er:YAG	2.94 μm	Free-running pulsed	Flexible fiber optic system or hollow waveguide; surface contact required for most procedures	Soft tissue incision and ablation; subgingival curettage; scaling of root surfaces; osteoplasty and ostectomy
Erbium, chromium:yttrium-selenium-gallium-garnet	Er,Cr:YSGG	2.78 μm	Free-running pulsed	Sapphire crystal inserts of varying diameters; surface contact required for most procedures	Soft tissue incision and ablation; subgingival curettage; osteoplasty and ostectomy
Neodymium:yttrium-aluminum-perovskite	Nd:YAP	1,340 nm	Pulsed	Flexible fiber optic system; surface contact required for most procedures	Soft tissue incision and ablation; subgingival curettage and bacterial elimination
Indium-gallium-arsenide-phosphide; gallium-aluminum-arsenide; gallium-arsenide	InGaAsP (diode) GaAlAs (diode) GaAs (diode)	Diodes can range from 635 to 950 nm	Gated or continuous	Flexible fiber optic system; surface contact required for most procedures	Soft tissue incision and ablation; subgingival curettage and bacterial elimination
Argon	Ar	488 to 514 nm	Gated or continuous	Flexible fiber optic system	Soft tissue incision and ablation

Oral Soft Tissues

Clinical applications. For many intraoral soft tissue surgical procedures, the laser is a viable alternative to the scalpel. In this regard, the literature is replete with numerous case reports and uncontrolled case studies reporting the use of various laser wavelengths, primarily CO₂, Nd:YAG, and diode, for intraoral soft tissue procedures, such as frenectomy, gingivectomy and gingivoplasty, deepithelization of reflected periodontal flaps, removal of granulation tissue, second-stage exposure of dental implants, lesion ablation, incisional and excisional biopsies of both benign

and malignant lesions, irradiation of aphthous ulcers, coagulation of free gingival graft donor sites, and gingival depigmentation.

Wound healing. The purported advantages of lasers versus scalpel surgery have been enumerated by numerous authors and include increased coagulation that yields a dry surgical field and better visualization; the ability to negotiate curvatures and folds within tissue contours; tissue surface sterilization and, therefore, reduction in bacteremia; decreased swelling, edema, and scarring; decreased pain; faster healing response; and increased patient acceptance.^{11,12}

Clearly, some of the claimed advantages are common sense conclusions based on clinical observations and patient response; for example, coagulation leading to better visualization of the surgical field and increased patient acceptance. Surprisingly, there are little data to support other contentions, such as faster healing response or decreased scarring. Indeed, claims of faster healing of laser soft tissue wounds appear to be wavelength specific and highly sensitive to energy density.

Most studies that examined healing rates of laser-induced wounds have involved the CO₂, Nd:YAG, or diode wavelengths. Studies concerning the CO₂ laser report that soft tissue healing is slower overall,¹³ slower initially but equal at 14 days,¹⁴ or equivalent¹⁵⁻¹⁷ compared to a conventional scalpel wound.

A comparison of wound healing following irradiation by the Nd:YAG and CO₂ lasers indicates that CO₂ laser-induced wounds in oral, oropharyngeal, and laryngeal mucosa healed significantly faster than those created by the Nd:YAG laser, but both heal slower than the conventional scalpel-induced wound.¹³ Delayed healing of Nd:YAG laser wounds compared to scalpel incisions also has been reported by Romanos et al.¹⁸ but only when using 3 W of power and a 20-Hz pulse rate. Healing was equivalent for scalpel and Nd:YAG wounds when the laser was used at a lower power setting of 1.75 W and 20 Hz.

Accelerated healing following laser-induced wound-ing has been reported but generally involves non-periodontal applications and use of soft lasers, e.g., low-level energy from a helium-neon diode.^{19,20} However, even this contention has been challenged by Damante et al.²¹ and Masse et al.,²² both of whom reported no evidence of accelerated healing following gingivectomy or periodontal flap surgery, using 670- and 810-nm diode lasers, respectively. Indirect evidence of accelerated healing from low-energy laser irradiation is offered by Crespi et al.,²³ who used the CO₂ laser in a defocused mode following surgical flap exposure of experimentally induced furcation involvements and reported induction of new periodontal ligament, cementum, and bone in Class III furcations in a dog experimental model. Lastly, almost as a casual observation, several studies have reported that laser-induced wounds show a decreased tendency toward scar contraction compared to traditional scalpel surgeries.^{24,25}

White et al.²⁶ used CO₂, Nd:YAG, Er:YAG, and two diode wavelengths (815 and 980 nm) to irradiate fresh bovine or porcine jaw tissue specimens. Based on tissue histology, it was determined that high power (watts), long pulse duration, high repetition rates (hertz), and long interaction times (duration of target exposure) all increased the risk of detrimental outcomes. In this

regard, it should be noted that numerous articles have been published in medical and veterinary journals comparing various laser wavelengths to electrosurgery and scalpel with regard to incisional time, blood loss, swelling and edema, pain, and general wound healing. However, the observations and conclusions of these studies are of limited value to clinical dentistry as most used power settings that ranged from five to 12 times greater than those used for dental surgical procedures. Furthermore, unlike oral mucosa, the target tissues in most of these studies were generally dermis or muscle with distant proximity to underlying bone.

Oral Hard Tissues

Effect of lasers on bone healing. Regardless of the type of instrumentation, the healing of bone following osteotomy, osteoplasty, or implant site preparation is complex, involving both local and systemic responses and a variety of cell types, enzymes, growth factors, cytokines, and other types of signal proteins. Exposure of bone to heating at levels $\geq 47^{\circ}\text{C}$ is reported to induce cellular damage leading to osseous resorption, and temperature levels of $\geq 60^{\circ}\text{C}$ result in tissue necrosis.²⁷ Given that laser/biologic tissue interactions are photothermal events that, in turn, are wavelength dependent, it should not be surprising that with the possible exception of two wavelengths (Er:YAG and Er,Cr:YSGG) the effect of most dental lasers on bone is generally detrimental.

There are relatively few studies measuring real-time bone surface temperatures while the overlying soft tissues are being irradiated by a dental laser. One such study by Fontana et al.²⁸ examined temperature increases at the bone surface while using an 810-nm diode laser within periodontal pockets in rats. Following 9 seconds of irradiation using 800 mW, 1.0 W, and 1.2 W of power delivered through a 300- μm optical fiber, they reported 10°C and 11°C increases in bone surface temperature. Only at a 600-mW setting was the bone surface temperature below the threshold for induction of cellular damage. If exposure time was lowered to 3 seconds, all power selections resulted in temperature increases that stayed below the critical threshold.

A second study²⁹ compared, *in vitro*, the CO₂ and Nd:YAG lasers using comparable energy densities for their effects on bone surface temperature while ablating overlying soft tissues. Energy densities ranged from 688 to 1286 J/cm², and all test runs were executed both with and without air/water surface cooling. Results showed that bone surface temperature increases ranged from 1.4°C to 2.1°C for the CO₂ laser and from 8.0°C to 11.1°C for the Nd:YAG laser. Although this was an *in vitro* study, these results indicated that when ablating relatively thin soft tissues supported by subjacent bone (e.g., mandibular facial

gingival and alveolar mucosa), the Nd:YAG laser should be used at low energy densities for short intervals; otherwise, there is a risk of irreversible bone damage.

Two studies have compared healing of tibial osteotomy defects in rats created by rotary bur, CO₂ (780 and 1,032 J/cm²) and Nd:YAG lasers (714 and 1,000 J/cm²). At all time intervals (0 to 63 days post-treatment), regardless of energy density or use of air/water surface coolant during irradiation, the osseous healing response was severely delayed.^{30,31}

Severe collateral damage has been identified as a major factor in delayed healing of laser-induced bone incisions. In studies reporting delayed healing, common observations appear to be the presence of a residual carbonized (char) layer on the treated surface, presence of inert bone fragments encapsulated by fibrous connective tissue, sequestra of bone, and bone fragments surrounded by multinucleated giant cells.^{17,30}

Given the limited collateral damage and efficacy of the Er:YAG laser when cutting enamel and root surfaces, one would expect numerous similar studies using bone as the target tissue. However, there are only about nine studies dating back to the mid-1980s. Two of the more recent studies employed a variety of techniques to evaluate the Er:YAG laser used at multiple energy settings, pulses/second (hertz), pulse durations, and water surface coolant to prepare osteotomy defects in a rat experimental model. Osteotomies created by the Er:YAG laser were compared to those created by rotary bur and the CO₂ laser.^{32,33} Collectively, the two studies indicate the Er:YAG laser, when used at a peak pulse energy of 100 mJ/pulse and 10 Hz, produced well-defined intrabony cuts with no evidence of melting or carbonization. Fourier transform infrared spectroscopy (FTIR), electron dispersive x-ray spectroscopy (EDX), and x-ray diffraction analysis revealed normal collagen/hydroxyapatite relationships covered by a thin surface layer characterized by a slight increase in calcium/phosphate ratio of the laser surface to be a result of formation of tetracalcium phosphate, which develops at temperatures above 1,100°C. Otherwise, the bone chemical composition was similar to that of defects produced by rotary bur. By contrast, CO₂ laser-induced osteotomies exhibited extensive charring, melting of the mineral phase, and delayed healing.

When considering laser-mediated osteotomy and/or ostectomy, the Er,Cr:YSGG appears to be a popular laser with clinicians. Yet, even with this wavelength, there is a paucity of evidence in the literature to support its use on bone. The clinical application of this wavelength to osseous resection and recontouring appears to be based on evidence derived from studies where the laser was used on enamel and dentin.

However, two recent studies suggest that the Er,Cr:YSGG wavelength may be suitable for use on bone. Kimura et al.³⁴ evaluated the morphological (scanning electron microscopy [SEM]), atomic (EDX), and temperature changes in canine mandibular bone, *in vitro*, following irradiation with the Er,Cr:YSGG laser at power settings of 5 W and 8-Hz pulse repetition for 10- or 30-second durations with concomitant air/water surface spray. The maximum temperature increase of 12.6°C was achieved only during the 30-second exposure. EDX analysis showed no change in calcium/phosphate ratio, and the SEM examination revealed cleanly cut bone with no evidence of charring or melting. Wang et al.³⁵ used the laser at an energy density of 80 J/cm² to create osseous perforations of 0.4-mm diameter in the maxilla and mandible of rabbits. They reported normal and complete healing of the wounds at 56 days post-treatment.

Laser-induced root surface modifications. Surface modification of cementum and dentin has been studied using a variety of laser wavelengths, primarily CO₂, Nd:YAG, Er:YAG, and, to a lesser extent, the diode laser.³⁶ A major conceptual consideration in laser-induced root surface modification is selection of a wavelength that will effectively remove calculus while suppressing both thermal damage to the pulp tissue and undesired removal of sound root structure. Achievement of these goals requires a wavelength characterized by minimal penetration depth in mineralized tissue. The mineral phase of both cementum and dentin is a carbonated hydroxyapatite that has intense absorption bands in the mid-infrared region. Consequently, of the laser wavelengths studied, the Er:YAG laser would appear to be the instrument of choice for effective removal of calculus, for root etching, and for creation of a biocompatible surface for cell or tissue reattachment. This latter statement is supported by Aoki et al.³⁶ in their definitive review of the literature concerning laser applications in non-surgical periodontal therapy.

In chronological terms, the CO₂ laser appears to have been the first of the currently available wavelengths to be studied for effects on root surfaces. Even at power settings as low as 4 W, the results of the earlier studies were not particularly encouraging, in that charring and melting of the root surface were common findings. In addition, FTIR analysis of charred surfaces revealed the presence of cyanamide and cyanate, both cytotoxic chemical residues.³⁷

More recent studies of the biocompatibility of CO₂ laser-treated surfaces, even when used at low energy densities, have yielded conflicting results. For example, both Pant et al.³⁸ and Crespi et al.³⁹ have reported increased *in vitro* attachment of fibroblasts to laser-treated surfaces compared to controls of SRP or chemically treated surfaces. On the other hand,

Fayad et al.⁴⁰ reported a total lack of fibroblast attachment to surfaces irradiated at only 1.25 mJ/pulse.

Gopin et al.⁴¹ used an *in vivo* animal model to demonstrate cytotoxicity of root surface char resulting from CO₂ laser irradiation subsequent to mucoperiosteal flap surgery. In all cases where root charring was observed, histologic examination revealed a lack of flap reattachment to the root surface. By contrast, all specimens treated by SRP alone or by laser irradiation followed by SRP exhibited flap reattachment to the treated root surfaces.

Heat-induced cracking of the root surface is a common observation when using the CO₂ laser, particularly at power settings of ≥ 4 W delivered in a continuous waveform. However, when used in a defocused mode, with pulsed waveform, and at low power settings, the CO₂ laser appears to inflict little damage.⁴² Indeed, using such parameters, the CO₂ laser has been shown to effectively remove smear layers.⁴³ Despite the latter results, due to the diameter of the hollow delivery tip (≥ 1 mm) that is required to transmit the energy beam, CO₂ lasers have restricted application in subgingival periodontal therapy.

Studies concerning applications of the Nd:YAG laser to root surfaces, such as those concerning the CO₂ laser, have yielded mixed results. Israel et al.⁴⁴ compared root surface changes following irradiation with CO₂, Nd:YAG, and Er:YAG lasers. At energy densities of 100 to 400 J/cm² for the CO₂ and 286 to 1,857 J/cm² for the Nd:YAG lasers, the authors noted that the degree of morphologic change following laser irradiation was directly related to energy density but unrelated to use of an air/water surface coolant. Changes in root surfaces included cavitation defects, globules of melted and resolidified mineral, surface crazing, and production of a superficial char layer.

Several *in vitro* studies^{36,45-47} have demonstrated heat-induced morphological changes of the root surface following irradiation with the Nd:YAG laser at power settings ranging from a low of 156.2 to 166.6 J/cm² to a high of 571 J/cm². As with the CO₂ laser, the Nd:YAG laser produced root surface alterations that included craters, charring, and melting and resolidification of the mineral phase. It should be noted, however, that even the lowest energy density used in these studies was still greater than those currently recommended for *in vivo* use. At least one study has reported altered chemical organization of root structure proteins following irradiation with the Nd:YAG laser at relatively low power settings (0.5 to 1.5 W).⁴⁸ Such heat-induced alterations in root structure proteins undoubtedly account for the separation of cementum from dentin following Nd:YAG irradiation and SRP, *in vitro*, reported by Morlock et al.⁴⁹

A recent study by Chen et al.,⁵⁰ in which cell cultures of human periodontal ligament fibroblasts were

subjected to Nd:YAG irradiation at low energy densities, reported significant decreases in cellular viability and collagen synthesis at 5 days post-treatment and evidence of mineralization of necrotic cells at 28 days post-treatment. Laser parameters were 50 mJ of power and 10 Hz, with a defocused beam delivered through a 400- μ m-diameter optical fiber, and durations of exposure ranging from 60 to 240 seconds.

In contrast with studies reporting detrimental results, at least two *in vitro* studies have demonstrated that the Nd:YAG laser, when used at low energy densities or a combination of low energy density with a defocused beam, can remove root surface smear layers without causing collateral damage to underlying cementum and/or dentin or increasing temperatures to a level that might trigger irreversible pulpal damage.^{51,52}

Two studies with similar experimental designs and diametric results demonstrate the possible effect that differing laser wavelengths can have on cells *in vitro*. The first study, using GaAs and GaAlAs diode lasers at energy densities between 0.95 and 6.32 J/cm², evaluated the effect of laser irradiation on prostaglandin E₂ (PGE₂) production and cyclooxygenase-1 (COX-1) and COX-2 gene expression in lipopolysaccharide-challenged human gingival fibroblasts.⁵³ The authors reported that irradiation with the GaAlAs diode laser significantly inhibited PGE₂ production in a dose-dependent manner, which, in turn, lead to reduction of COX-2 mRNA levels. The second study irradiated cell cultures of human gingival fibroblasts with an Er:YAG laser at energy densities ranging from 1.68 to 3.37 J/cm² and actually increased production of PGE₂ and COX-2 mRNA.⁵⁴

Other *in vitro* studies have reported a lack of positive effect on attachment of periodontal ligament cells to root surfaces following low-level irradiation (1 W for 20 seconds) with the GaAlAs diode laser⁵⁵ or detrimental ultrastructural changes that could potentially lead to disturbances in collagen synthesis.⁵⁶ Despite these reports, Kreisler et al.⁵⁷ reported that low-level irradiation with the GaAlAs diode laser (10 mW for 75, 150, and 300 seconds) had a stimulatory effect on the proliferation of periodontal ligament fibroblasts *in vitro*. Given the findings of these studies, one might conclude that differing levels of power and times of exposure produce different interactive results that, in turn, indicate an irradiation threshold above which cell damage is likely to occur.

Another example of the importance of parameter selection was reported by Kreisler et al.⁵⁸ following *in vitro* irradiation of tooth root specimens with the GaAlAs diode. The study reported little to no damage on the root surface at a power output of ≤ 1 W, whereas power selections of 1.5, 2.0, and 2.5 W produced varying degrees of carbonization (charring) and heat-induced surface cracking.

Due to its high absorption in both water and hydroxyapatite, the bulk of recent research concerning laser-induced root surface modification has involved the Er:YAG laser. This wavelength of laser has been shown to effectively remove smear layers,^{44,59} dental calculus,⁶⁰⁻⁶³ cementum,^{61,64} and cementum-bound endotoxin.⁶⁵ When used at low energy densities with a water spray surface coolant, the majority of studies report little to no heat-induced tissue damage and production of smooth root surfaces.^{65,66} In addition, *in vitro* fibroblast adhesion studies show that the resultant root surface appears to be at least as biocompatible as that produced by SRP.^{60,67,68}

There are, however, caveats with respect to the Er:YAG laser and root surface interactions. First, the water spray surface coolant appears important in suppressing heat-induced surface alterations and protection of the pulp against elevations in temperatures during root surface irradiation.^{63,64,66,69} Achievement of adequate water cooling in deeper periodontal pockets is likely to be inconsistent. Second, most studies reporting heat-induced surface damage or lack thereof have used SEM. Obviously, the SEM is a surface-scanning microscope and does not detect subsurface damage. At least one study has noted subsurface alterations in dentin following ablation of cementum by the Er:YAG laser using power settings of 60, 100, and 180 mJ. The thermal changes extended into the dentin from 255 to 611 μm , as measured from the target surface, and appeared independent of irradiation energy.⁷⁰ Third, as with other laser wavelengths, the selection of parameters is of paramount importance when discussing tissue damage. In this regard, Crespi et al.⁷¹ in an *in vitro* study noted that use of the Er:YAG laser in a defocused non-contact mode effectively removed calculus with only minimal removal of cementum. Several studies emphasize the relationship between increasing power and energy density and increased removal of root surface structure^{63,64} and the number and depth of craters in the target surface.^{61,64}

A relatively new laser, the Nd:YAP with a wavelength of 1,340 nm, has been tested on the root surfaces of extracted teeth. The authors reported the presence of heat-induced damage at energy densities ranging from 509 to 1,274 J/cm². However, as might be expected, the degree of damage was directly related to increasing energy density and progressively evolved from simple surface cracking of cementum to deep cratering to melting and deep ablation of cementum with exposure of the underlying dentin.⁷²

Use of lasers for clinical crown lengthening without gingival flap reflection. Recently, the Er,Cr:YSGG laser, and to a lesser extent the Er:YAG laser, has been promoted for clinical crown lengthening without gingival flap reflection for both esthetic and prosthetic rea-

sons.⁷³⁻⁷⁸ These articles generally fall into one of two categories: non-controlled case studies and technique-oriented articles. Collectively, the articles raise several questions: 1) Is there sufficient tactile sensation transmitted through the laser delivery tip to allow the clinician to adequately distinguish between bone and root surface cementum and/or dentin? 2) Have any of these reports determined if the roots of treated teeth incur surface damage, e.g., cratering, ditching, charring, heat-induced cracking, or melting? 3) In cases requiring bone removal, does the lack of direct visualization allow the clinician to establish proper anatomical dimensions and contours that will maintain the gingival papilla post-surgically and prevent violation of the biologic width?

Currently, there are no controlled longitudinal or cohort studies supporting use of lasers for clinical crown lengthening using the closed-flap technique. Thus, there are no satisfactory answers to any of these questions because there are no published research data. The only existing support for such applications are non-controlled case reports. Obviously, esthetic crown lengthening can easily be managed with lasers if clinically short crowns are the result of gingival overgrowth or lack of passive eruption. However, in such cases, there is increased probing depth (PD) due to excessive amounts of soft tissue, and violation of the biologic width is usually not a major concern.

Effect of lasers on bacteria and calculus. The use of a dental laser in the treatment of chronic periodontitis is based on the purported benefits of subgingival curettage, laser-induced new attachment through regeneration of cementum, periodontal ligament, and supporting alveolar bone, and significant decreases in subgingival pathogenic bacteria. There is limited evidence suggesting that lasers effect greater reductions in subgingival bacteria than that achieved by traditional therapy. Indeed, most laser bactericidal studies have been *in vitro* investigations that have little relevance to the protected biofilms of a periodontal pocket. Indeed, regarding subgingival biofilms, such factors as thickness of microbial mass, density and thickness of extracellular matrix, microbial composition, color, and water content have yet to be replicated *in vitro*. Most laser bactericidal studies report a dose/response relationship; that is, increases in power or energy density result in increased destruction of bacteria. However, in many studies, energy densities are often not reported or cannot be calculated due to incomplete listing of parameters. Studies also vary in how the laser energy is delivered to the target surface, some using a sweeping motion of the delivery tip and others using a static exposure of single or multiple pulses. Lastly, the angle of irradiation can vary from 0 to 90°, making computation of energy densities nearly impossible. Despite these problems, one can

still discern trends in the literature regarding the bactericidal effects of dental lasers.

One of the first in vivo studies reporting reductions in pathogenic bacteria following irradiation with the Nd:YAG laser showed decreases in *Porphyromonas gingivalis* (*Pg*), *Prevotella intermedia* (*Pi*), and *Actinobacillus actinomycetemcomitans* (*Aa*). However, teeth extracted 7 days post-treatment exhibited recolonization of laser-irradiated subgingival root surfaces by multiple morphotypes of bacteria.⁷⁹

A later study,⁸⁰ also using the Nd:YAG laser, compared laser therapy to SRP and reported that both modalities reduced levels of *Tannerella forsythensis* (*Tf*), *Pg*, and *Treponema denticola* (*Td*) but incompletely eliminated *Aa*. Laser therapy resulted in a greater reduction in microbial levels than did SRP, although both treatments exhibited microbial rebound approaching baseline levels at 10 weeks post-therapy.

A third in vivo study compared SRP (one episode) to SRP followed by irradiation with the Nd:YAG laser at a relatively high energy density of 124 J/cm². Treated pockets were irradiated once per week for 3 weeks. Levels of *Pg*, *Pi*, and *Aa* were determined at 6 months post-treatment, and only levels of *Pg* were found to be significantly reduced compared to SRP.⁸¹

In vitro studies using the Nd:YAG laser at low power settings have reported calculus ablation without detrimental effects to underlying cementum or dentin;⁸² a linear relationship between energy level, microbial numbers, and concentration of hemoglobin (blood) and minimal energy required for a bactericidal effect;⁸³ a differential susceptibility of various microbes to laser energy;⁸³ and a differential susceptibility to damage of calculus, cementum, and dentin, even within the same specimen.⁸² The latter phenomenon is likely the result of such factors as variability in color, thickness, composition, texture, and water content.

There are relatively few articles regarding diode lasers and their respective interactions with bacteria and dental calculus. A well-designed study by Harris and Yessik⁸⁴ determined the in vitro ablation threshold for *Pg* for both the 810-nm diode and Nd:YAG lasers to be 48 and 96 J/cm², respectively. The diode laser (805 nm), when used adjunctively with SRP, has been shown to have an additive effect in reducing subgingival bacterial populations in periodontal pockets of ≥ 4 mm in depth.⁸⁵ Interestingly, the diode laser used at low intensity (1 J/cm²) has also been shown to have an in vitro stimulatory effect on bacterial growth that appears to be species specific.⁸⁶

Coffelt et al.⁸⁷ demonstrated in vitro bacterial ablation with the CO₂ laser in a defocused mode at an energy density of 11 J/cm². They also determined the threshold energy density for inducing root damage to be 41 J/cm², well above that required to destroy bacteria adherent to a root surface. Following this line

of thought, Crespi et al.⁸⁸ demonstrated that the use of SRP followed by CO₂ laser irradiation at an energy density of 2.45 J/cm² produced root surfaces devoid of residual bacteria. One caveat, however, is that ablation of bacteria requires a direct hit by the energy beam. There appears to be a well-defined interface between irradiated bacteria and those not damaged by the energy beam.⁸⁹ This factor may explain why many in vivo studies show the persistence of viable bacteria following subgingival laser irradiation.

As with other wavelengths, studies involving the Er:YAG laser are predominantly in vitro investigations. Collectively, these studies report bacterial ablation at energy densities as low as 0.3 J/cm²,⁹⁰ effective removal of calculus without associated heat-related damage to the root structure at energy densities of ≤ 10.6 J/cm²,^{91,92} and removal of root structure without significant increases in pulp chamber temperatures.⁹¹

By contrast, Aoki et al.⁹³ reported an equivalent effect when using ultrasonic instrumentation or the Er:YAG laser at an energy density of 14.2 J/cm² for the in vitro removal of calculus from extracted teeth. The laser produced slightly rougher topography and thermal microchanges on the root surface. Folwaczny et al.⁹⁴ offer another caveat, in that in vitro use of the Er:YAG laser (60 mJ, 15 Hz, 250-microsecond pulse duration, and a total of 55, 75, or 105 pulses) on root surfaces seeded with bacteria only reduced bacterial loads by one log. Lastly, Eberhard et al.⁹⁵ compared laser removal of calculus to SRP in situ. Microbial samples were taken prior to and immediately after treatment for DNA probe analysis. Following tooth extraction, SEM using digitized planimetry was used to measure residual calculus. Results showed that only 68.4% of the root surface was calculus free in contrast to 94% after SRP. If the laser was used for twice the time as that for SRP, the percentage of root surface devoid of calculus increased to 83.3%. Both treatments resulted in similar reductions of pathogenic microbes.

Treatment of Chronic Periodontitis

As previously stated, laser-mediated periodontal therapy is predicated on the concept of subgingival curettage and/or reattachment and regeneration of the attachment apparatus. Such laser therapy is commonly referred to as “non-surgical.” Clearly, use of the term non-surgical when referring to a procedure based on the concept of subgingival curettage is debatable. Be that as it may, there are no convincing data that a regenerated connective tissue attachment is superior to attachment via a long-junctional epithelium, the latter commonly a result of non-surgical mechanical therapy.^{96,97} The desirability of the connective tissue attachment is based on hypothesis, not fact. Indeed, at least two investigations report that attachment mediated by long-junctional epithelium may

Table 4.**Laser Treatment of Chronic Periodontitis: Summary of Longitudinal Clinical Trials and Cohort Studies**

Study (listed in chronological sequence)	Type of Laser	Number of Patients	Length of Study (days)	Initial PD (mm)	Decrease in Subgingival Bacteria: Laser Versus Control
Rydén et al. ¹¹²	GaAs diode	10	28	NA (gingivitis study)	No statistical difference between treatment groups.
Finkbeiner ¹¹⁶	Argon	30	138	4 to 5 6 to 7 8 to 9	NA
Ben Hatit et al. ⁸⁰	Nd:YAG	14	70	≥5	No significant difference. Reported reductions in <i>Tf</i> , <i>Pg</i> , and <i>Td</i> but not <i>Aa</i> . All microbes rebounded to baseline at 10 weeks.
Radvar et al. ¹¹⁷	Nd:YAG	11	42	>4	At 6 weeks, only the SRP group showed a significant reduction from baseline.
Neill and Mellonig ¹⁰¹	Nd:YAG	10	180	>4	No significant difference for <i>Pg</i> or <i>Pi</i> .
Moritz et al. ¹¹⁵	GaAs diode	50	180	3.9 versus 3.0 (mean depth in molar region)	No significant difference. 59% of lased sites had 1 log decrease versus 33% of controls; 27% of lased sites had 2 log decrease versus 17% of controls.
Liu et al. ¹¹⁸	Nd:YAG	8	84	4 to 6	NA
Schwarz et al. ¹⁰³	Er:YAG	20	180	≥4	No significant difference between treatment groups.
Gutknecht et al. ⁸¹	Nd:YAG	20	175	4 to 6	No significant difference between treatment groups.

Table 4. (continued)**Laser Treatment of Chronic Periodontitis: Summary of Longitudinal Clinical Trials and Cohort Studies**

Mean Reduction in PD: Laser Versus Control (mm)	Mean Gain in CAL: Laser Versus Control (mm)	Percentage of Decrease in BOP: Laser Versus Control	Comments
NA	NA	No statistical difference between test and controls.	Total dose of 1 J/cm ² over 4 minutes; laser did not influence the inflammatory reaction of the marginal gingiva in an experimental human gingivitis model. Controls were untreated.
1.62 versus NA 2.85 versus NA 3.30 versus NA	NA	75% versus NA	Used 0.4 W and 0.3-mm-diameter optical fiber with a 20- to 30-second exposure; no control group.
NA	NA	NA	Treatment groups: 0.8 W at 10 Hz and 100 mJ/pulse versus 1.0 W at 10 Hz and 100 mJ/pulse versus 1.2 W at 12 Hz and 100 mJ/pulse versus 1.5 W at 15 Hz and 100 mJ/pulse versus scaling only (control); pulse duration was 150 microseconds, and laser was fitted with 0.3-mm-diameter optical fiber.
0.50 versus 1.70	NA	<10% versus 45%	Treatment groups: 50 mJ versus 80 mJ (test) versus SRP (control); pulse duration of 150 microseconds using a 0.32-mm-diameter optical fiber; energy densities were 62.9 and 99.5 J/cm ² .
1.70 versus 0.50	1.1 versus 1.0	Significant difference but no data presented.	Treatment groups: SRP alone versus laser + SRP versus untreated control; 80 mJ at 25 Hz for 2 minutes; <4-mm pocket lased for 5 to 10 seconds; 4 to 6 mm lased for 20 seconds; 7 to 9 mm lased for 30 seconds; and >9 mm lased for 40 seconds.
1.30 versus 0.40	NA	Study used papillary bleeding index: improvement in 97% of lased versus 67% of controls.	Used 2.5 W and 50 Hz, and 10-microsecond pulse duration; controls were scaling followed by H ₂ O ₂ rinsing at 1 week and at 2 and 4 months. Tests were scaled plus lased at 1 week and at 2 and 4 months.
NA	NA	NA	Laser versus SRP versus laser + SRP versus SRP + laser; 150 mJ at 20 pulses per second; determined that SRP was required to reduce levels of gingival crevicular fluid interleukin-1 β .
2.00 versus 1.60	1.9 versus 1.0	77% versus 56%	Laser used at 160 mJ/pulse at 10 Hz (test) versus SRP (control).
0.85 versus 0.80 (estimated from line graph)	NA	85% versus 75% (estimated from line graph)	Used 100 mJ and 20 Hz, 100-microsecond pulse duration, 0.32-mm-diameter optical fiber, and energy density of 124 J/cm ² at fiber tip. Control was untreated versus one-time SRP versus SRP + laser treatment of pocket once per week for 3 weeks. Measured levels of Pg, Pi, and Aa. Found no significant difference for Pi and Aa; difference for Pg was significant.

Table 4. (continued)**Laser Treatment of Chronic Periodontitis: Summary of Longitudinal Clinical Trials and Cohort Studies**

Study (listed in chronological sequence)	Type of Laser	Number of Patients	Length of Study (days)	Initial PD (mm)	Decrease in Subgingival Bacteria: Laser Versus Control
Sjöstrom and Friskopp ¹¹⁹	Nd:YCG (1,061 nm versus 1,064 nm for Nd:YAG)	27	120	≥4	NA
Yilmaz et al. ¹¹³	GaAs diode	10	32	4	No significant difference between treatment groups.
Miyazaki et al. ¹⁰²	Nd:YAG	18	84	≥5	NA
Schwarz et al. ¹⁰⁴	Er:YAG	20	365	≥4	No significant difference between treatment groups.
Schwarz et al. ¹⁰⁵	Er:YAG	20	2 years	≥4	No significant difference between treatment groups.
Borrajó et al. ¹⁰⁰	InGaAsP diode	30	42	NA	NA
El Yazami et al. ¹⁰⁹	Nd:YAP	22	180	5.5 test, 5.2 control	NA
Harris et al. ¹²⁰	Nd:YAG	75*	180	4 to 6 7 to 9	NA
Sculean et al. ¹⁰⁶	Er:YAG	20	180	≥4	NA
Schwarz et al. ¹⁰⁷	Er:YAG	22	180	8.6	NA
Sculean et al. ¹⁰⁸	Er:YAG	23	180	7.8	NA
Ambrosini et al. ¹¹⁰	Nd:YAP	30	90	4.1	No significant difference between treatment groups.

Table 4. (continued)**Laser Treatment of Chronic Periodontitis: Summary of Longitudinal Clinical Trials and Cohort Studies**

Mean Reduction in PD: Laser Versus Control (mm)	Mean Gain in CAL: Laser Versus Control (mm)	Percentage of Decrease in BOP: Laser Versus Control	Comments
1.4 versus 1.4	NA	Gingival bleeding index showed no difference between treatment groups.	Treatment groups: laser + SRP + laser (test) versus SRP only (control); laser used at 7 W and 20 Hz with pulse length of 250 microseconds.
0.23 for laser only; 0.49 for SRP only; 0.66 for laser + SRP	NA	17% for laser only; 50% for SRP only; 60% for laser + SRP	Treatment groups: laser only versus laser + SRP versus SRP alone versus oral hygiene instructions alone; laser used at power density of 1.6 J/cm ² on days 1, 2, 4, 7, 9, and 11 using methylene blue dye as a photosensitizer.
1.43 for Nd:YAG; 1.36 for scaling; 1.00 for CO ₂	0.50 for Nd:YAG; 0.57 for scaling; 0.31 for CO ₂	43% for Nd:YAG; 34% for scaling; 16% for CO ₂	Treatment groups: Nd:YAG laser alone versus CO ₂ alone versus ultrasonic scaling alone; Nd:YAG laser used at 100 mJ/pulse and 20 Hz for 2 minutes; CO ₂ laser used at 2 W for 2 minutes.
2.0 for laser + SRP; 1.7 for laser only	Both treatments had 1.6-mm gains in CAL.	14% for laser + SRP; 16% for laser only	Treatment groups: laser + SRP (test) versus laser only (control); laser used at 160 mJ/pulse at 10 Hz.
1.60 versus 1.30	1.40 versus 0.70	64.3% versus 46.2%	Laser used at 160 mJ/pulse at 10 Hz versus SRP; this article reports the long-term results of the Schwarz et al. ¹⁰³ study.
NA	0.81 versus 0.85	72% versus 53%	SRP versus SRP + laser; 2 W, 100-millisecond pulse length, 50 Hz, 2-mm-diameter optical fiber.
2.80 versus 1.30	2.60 versus 1.10	67.1% versus 51.2%	70 mJ and 30 Hz; SRP (control) versus SRP + laser (test).
1.55 versus NA 3.44 versus NA	NA	NA	Laser power ranged from 3.0 to 4.8 W depending on operator; 1 minute/tooth exposure delivered a total energy of 1 to 15 J/mm of PD. LANAP protocol. Study also includes data from the Neill and Mellonig ¹⁰¹ study. Controls were historic, i.e., compared to other studies.
1.52 versus 1.57	1.11 versus 1.11	23% versus 31%	Laser used at 160 mJ/pulse and 10 Hz (test) versus ultrasonic scaling (control).
4.00 versus 4.1	3.2 versus 3.3	35% versus 26%	Access flap surgery + laser debridement + enamel matrix protein derivative (test) versus access flap surgery + SRP + enamel matrix protein derivative (control); laser used at 160 mJ/pulse and 10 Hz.
3.7 versus 3.2	2.6 versus 1.5	62.5% versus 59%	Laser used at 160 mJ/pulse and 10 Hz (test) versus flap surgery and debridement of root and defect (control).
1.50 versus 1.30	1.00 versus 1.10	85.7% versus 85.7%	SRP only (control) versus SRP + laser (testy); bacteria tested by DNA probe: Aa, Pi, Pg, Tf, and Td; laser parameters were 10 W and 0.2-mm-diameter optical fiber.

Table 4. (continued)**Laser Treatment of Chronic Periodontitis: Summary of Longitudinal Clinical Trials and Cohort Studies**

Study (listed in chronological sequence)	Type of Laser	Number of Patients	Length of Study (days)	Initial PD (mm)	Decrease in Subgingival Bacteria: Laser Versus Control
Noguchi et al. ¹¹¹	Nd:YAG	16	90	5.8 for control; 4.9 for laser; 5.2 for laser + minocycline; 5.4 for laser + irrigation	Proportions of <i>Pg</i> , <i>Pi</i> , and <i>Tf</i> were significantly decreased in laser + minocycline group compared to laser only and laser + irrigation groups.
Qadri et al. ¹¹⁴	InGaAlP and GaAlAs diodes	17	42	4.7	No significant difference between treatment groups.

NA = not applicable because study did not measure parameter.

* Includes 10 patients from Neill and Mellonig¹⁰¹ study.

be as resistant to disease as a true connective tissue attachment.^{98,99}

A meaningful comparison between various clinical studies or between laser and conventional therapy is difficult at best and likely impossible at the present. Reasons for this dilemma are several, such as different laser wavelengths; wide variations in laser parameters; insufficient reporting of parameters that, in turn, does not allow calculation of energy density; differences in experimental design; lack of proper controls; differences in severity of disease and treatment protocol; and measurement of different clinical endpoints. Despite these problems and given that, to date, only 23 human clinical trials have been published,^{80,81,100-120} one may still extract sufficient data to recognize trends in the results of laser-mediated treatment of chronic periodontitis (Table 4).

When measuring outcomes of non-surgical periodontal therapy, gain in clinical attachment level (CAL) represents the gold standard. PD and levels of subgingival microbes are important primarily because, in cases of traditional non-surgical mechanical therapy, they have been shown to be associated with changes in CAL.¹⁰ Thus, it is somewhat surprising that only 12 of the 23 laser clinical trials considered gains in CAL as an endpoint of treatment (Table 4). If one calculates the average gain in CAL reported in 11¹⁰⁰⁻¹¹⁰ of the 12 studies for laser therapy versus con-

trols (1.62 mm versus 1.26 mm, respectively), it becomes evident that there is minimal benefit following subgingival laser therapy. One study¹¹¹ was not included due to use of local drug delivery combined with laser therapy and lack of an SRP control group. Further analysis of these data shows that seven of the studies^{100-102,104,106,107,110} reported essentially equivalent results between laser-treated groups and controls, and four studies^{103,105,106,109} clearly favored laser therapy (average gain of 2.13 mm in CAL following laser therapy versus 1.05 mm for controls). Of the latter four studies, three involved the Er:YAG laser^{103,105,108} and one the Nd:YAP laser,¹⁰⁹ and control groups consisted of SRP (three studies) and surgical flap with debridement (one study).¹⁰⁷

There are presently five peer-reviewed published clinical trials using a diode laser wavelength. Two of these are actually an investigation of the effects of low-level laser therapy (LLLT) on the gingival inflammatory response in a human experimental gingivitis model.^{99,111} Results showed that LLLT had no impact on bacterial plaque accumulation, bleeding on probing (BOP), or the number of vessels in the marginal gingiva, all being indicators of the inflammatory response.

Of the three remaining studies, three essentially reported little to no effect on bacterial levels¹¹³⁻¹¹⁵ compared to controls and minimal changes in both PDs^{113,114} and BOP.^{96,109} Only one article has reported

Table 4. (continued)**Laser Treatment of Chronic Periodontitis: Summary of Longitudinal Clinical Trials and Cohort Studies**

Mean Reduction in PD: Laser Versus Control (mm)	Mean Gain in CAL: Laser Versus Control (mm)	Percentage of Decrease in BOP: Laser Versus Control	Comments
1.57 for laser; 2.39 for laser + minocycline; 1.60 for laser + irrigation; no change for control	1.52 for laser; 2.36 for laser + minocycline; 1.62 for laser + irrigation; no change for control	63% for laser; 77% for laser + minocycline; 65% for laser + irrigation; 19% for control	Used 200 mJ and 10 Hz for 90 seconds with 0.4-mm-diameter optical fiber; treatment groups were laser only, laser + local minocycline, laser + povidone-iodine irrigation, and sham procedure (control). Each treatment was done every week for 1 month.
0.90 versus 0.20	NA	NA	First used InGaAlP laser with energy density of 4.5 J/cm ² and then the GaAlAs laser in pockets ≥6 mm at energy density of 8.75 J/cm ² . Control was laser placebo (sham treatment with laser inactivated); measured levels of gingival crevicular fluid elastase, interleukin-1β, and matrix metalloproteinase-8 and found no significant differences resulting from laser treatment.

NA = not applicable because study did not measure parameter.

* Includes 10 patients from Neill and Mellonig¹⁰¹ study.

significant effects on clinical parameters that favored the diode laser wavelength.¹¹⁵ However, this study used inappropriate controls for comparison, i.e., both test and control patients received scaling of periodontally diseased sites, but sites in test patients were repeatedly lased at 1 week and 2 and 4 months, while control patients received no further treatment except to rinse with H₂O₂.

Finkbeiner¹¹⁶ in an uncontrolled cohort study reported the effects of treating chronic periodontitis using an argon laser combined with subgingival scaling and chlorhexidine irrigation. Results showed reductions of 1.62, 2.85, and 3.30 mm for 4- to 5-, 6- to 7-, and 8- to 9-mm initial PDs, respectively. In addition, a 75% decrease in BOP was noted.

At present, there are only nine published clinical trials using the Nd:YAG laser for treatment of chronic periodontitis. When viewed as a collective body of evidence, they provide conflicting results.^{80,81,101,102,111,117-119} For example, two studies did not measure PD as an endpoint,^{80,118} three studies reported little to no difference in PD reduction when comparing laser-treated sites to control sites (SRP only),^{81,102,109} and one study reported a greater mean decrease in PD in SRP-treated sites than in those treated by laser.¹¹⁷ Of the remaining three studies, the laser improved PD compared to untreated controls¹⁰¹ or historic controls (i.e., data reported in other studies

used for comparison).¹²⁰ However, the study using historic controls¹²⁰ reported relatively large standard deviations for mean PD reductions in laser-treated pockets, indicating a variation in technique or a degree of unpredictability. Lastly, when the Nd:YAG laser was combined with locally delivered minocycline, PD was significantly reduced compared to sites treated by laser alone.¹¹¹

In this same series of studies, results for BOP and reduction of pathogenic bacteria followed the same general trend as PD reduction. Take, for example, reductions in specific periodontal pathogens: four studies did not measure this factor as an endpoint,^{102,118-120} two studies reported no difference between test and control groups in levels of subgingival bacteria,^{81,101} one study reported that SRP was more effective than laser therapy,¹¹⁷ and two studies reported significant decreases in laser-treated sites of *Tf*, *Td*, *Pg*, or *Pi*.^{80,111} Interestingly, one study reported significant decreases in subgingival bacterial levels for both Nd:YAG laser-treated sites and SRP, but there was no difference between treatment groups in levels of *Pg* or *Pi*.¹⁰¹ Lastly, one study reported that SRP was necessary to effect significant decreases in gingival crevicular fluid levels of interleukin-1β compared to laser only or combinations of laser and SRP.¹¹⁸

In general, clinical trials involving the Er:YAG laser in the treatment of chronic periodontitis are better

designed and yield more consistent results. A major reason for this distinction is likely related to the fact that all six clinical trials have come from the same group of investigators.¹⁰³⁻¹⁰⁸ Four of the six studies compared the effects of the Er:YAG laser alone or in combination with SRP to SRP alone.¹⁰³⁻¹⁰⁶ The remaining two studies combined the laser with periodontal surgery or enamel matrix protein in the treatment of intrabony defects.^{107,108}

With respect to reductions in subgingival bacterial loads, comparisons of Er:YAG laser therapy (alone or as an adjunct to SRP) to SRP alone were reported to have no significant difference between treatments, i.e., both groups showed a significant increase in cocci and non-motile rods and a decrease in the numbers of motile rods and spirochetes.¹⁰³⁻¹⁰⁵ Decreases in mean PD consistently favored the laser treatment, but differences between treatment groups ranged from essentially no difference^{107,108} to 0.4 at 6 months⁹⁸ and 1 year¹⁰⁵ to 0.3 mm at 1 year¹⁰⁴ and 2 years post-treatment.¹⁰⁵ Lastly, percentage of reduction in BOP in these studies was somewhat conflicted in that three studies favored the laser versus SRP,^{103,105,107} two studies reported essentially equivalent reductions in BOP (14% versus 16% for laser + SRP versus laser alone¹⁰⁴ and 62.5% versus 59% for flap surgery + laser versus flap surgery + traditional debridement),¹⁰⁸ and one study favored SRP alone versus laser alone.¹⁰⁶ None of the BOP comparisons were reported as statistically significant differences.

A relatively new laser wavelength to dentistry is Nd:YAP (1,340 nm), which has an absorption coefficient in water approximately 20 times greater than the Nd:YAG laser (1,060 nm). Thus far, only two clinical trials using this laser in the treatment of chronic periodontitis have been published,^{109,110} both comparing SRP + laser to SRP alone.

One study¹⁰⁹ reported a 1.5-mm mean difference in PD reduction (initial mean PD \approx 5.5 mm) that favored adjunctive use of the laser with SRP versus SRP alone. In addition, there was a reported difference in reduction of BOP between the two treatment groups of 15%, again favoring the combined treatment protocol. The second study¹¹⁰ measured plaque index, gingival index, BOP, PD, CALs, and the presence of *Aa*, *Pg*, *Pi*, *Tf*, and *Td*. The authors reported no significant difference between treatment groups for any of the measured clinical parameters, i.e., no additional advantage was achieved by using the Nd:YAP laser.

Examination of all 23 clinical trials reveals that blinded examiners were reported as part of the experimental design in only four of the studies,^{100,104,109,110} and calibration of examiners was reported in six studies.¹⁰⁴⁻¹⁰⁹ Such oversights in experimental design raise the issue of introducing bias when reporting experimental results. It is well accepted that the ultimate

applicability and strength of recommendation for a specific treatment modality must be based on the weight of evidence. In this regard, a single randomized, blinded, controlled, longitudinal, clinical trial carries more weight than even a large series of case observations because of the greater potential for bias when observations are made under uncontrolled conditions. Although the dental literature is replete with clinical observations that have evolved from private practice settings, such studies rarely present concurrent controls; therefore, effectiveness of treatment is likely to be overestimated.

CONCLUSIONS

The results achieved following irradiation of biologic tissue by a specific wavelength of laser is directly related to the selected parameters. In other words, given the same wavelength, different laser parameters will yield different levels of energy density for varying periods of time and, thereby, different degrees of change in the target tissue. This review consistently noted that even when using the same wavelength of laser, there was little concurrence in the choice of parameters in the experimental methods. This recurring problem makes comparison of results nearly impossible and undoubtedly accounts for many of the reported conflicts in results for the various laboratory studies and clinical trials.

Based on this review of the literature, one must conclude that there is a great need to develop an evidence-based approach to the use of lasers for the treatment of chronic periodontitis. Simply put, there is insufficient evidence to suggest that any specific wavelength of laser is superior to the traditional modalities of therapy. Current evidence does suggest that use of the Nd:YAG or Er:YAG wavelengths for treatment of chronic periodontitis may be equivalent to SRP with respect to reduction in PD and subgingival bacterial populations. However, if gain in CAL is considered the gold standard for non-surgical periodontal therapy, then the evidence supporting laser-mediated periodontal treatment over traditional therapy is minimal at best. Lastly, there is limited evidence suggesting that lasers used in an adjunctive capacity to SRP may provide some additional benefit. Establishment of a sound evidence base for laser usage in treatment of chronic periodontitis will require randomized, blinded, controlled, longitudinal, clinical trials. Given the inherent expense, requirements of time, and number of clinicians required to conduct such studies, this may require multicenter collaborative studies.

ACKNOWLEDGMENT

The author thanks the staff of the University of Missouri School of Dentistry library for their considerable

effort in locating and copying the substantial number of articles reviewed for this article.

REFERENCES

- Maiman TH. Stimulated optical radiation in ruby. *Nature* 1960;187:493-494.
- Snitzer E. Optical maser action of Nd⁺³ in a barium crown glass. *Phys Rev Lett* 1961;7:444-446.
- Goldman L, Hornby P, Meyer R, Goldman B. Impact of the laser on dental caries. *Nature* 1964;203:417.
- Stern RH, Sognaes RF. Laser inhibition of dental caries suggested by first tests in vivo. *J Am Dent Assoc* 1972;85:1087-1090.
- Myers TD, Myers WD. In vivo caries removal utilizing the YAG laser. *J Mich Dent Assoc* 1985;67:66-69.
- Myers TD. What lasers can do for dentistry and you. *Dent Manage* 1989;29:26-28.
- Midda M, Renton-Harper P. Lasers in dentistry. *Br Dent J* 1991;170:343-346.
- Midda M. Lasers in periodontics. *Periodontol Clin Investig* 1992;14:14-20.
- Midda M. The use of lasers in periodontology. *Curr Opin Dent* 1992;2:104-108.
- Cobb CM. Non-surgical pocket therapy: Mechanical. *Ann Periodontol* 1996;1:443-490.
- Wigdor H, Walsh J, Featherstone JDB, et al. Lasers in dentistry. *Lasers Surg Med* 1995;16:103-133.
- Bader H. Use of lasers in periodontics. *Dent Clin North Am* 2000;44:779-792.
- Lippert BM, Teymoortash A, Folz BJ, Werner JA. Wound healing after laser treatment of oral and oropharyngeal cancer. *Lasers Med Sci* 2003;18:36-42.
- Romanos G, Chong HS, Ng K, Toh CG. A preliminary study of healing of superpulsed carbon dioxide laser incisions in the hard palate of monkeys. *Lasers Surg Med* 1999;24:368-374.
- Arashiro DS, Rapley JW, Cobb CM, Killoy WJ. Histologic evaluation of porcine skin incisions produced by CO₂ laser, electrosurgery and scalpel. *Int J Periodontics Restorative Dent* 1996;16:479-491.
- Arcoria CJ, Steele RE, Vitasek BA, Wagner MJ. Effects of coaxial CO₂/Nd:YAG irradiation on periodontal wound healing. *Lasers Surg Med* 1992;12:401-409.
- Williams TM, Cobb CM, Rapley JW, Killoy WJ. Histologic evaluation of alveolar bone following CO₂ laser removal of connective tissue from periodontal defects. *Int J Periodontics Restorative Dent* 1995;15:497-506.
- Romanos GE, Pelekanos S, Strub JR. A comparative histologic study of wound healing following Nd:YAG laser with different energy parameters and conventional surgical incision in rat skin. *J Clin Laser Med Surg* 1995;13:11-16.
- Fernando S, Hill CM, Walker R. A randomized double blind comparative study of low level laser therapy following surgical extraction of lower third molar teeth. *Br J Oral Maxillofac Surg* 1993;31:170-172.
- Neiburger EJ. The effect of low-power lasers on intra-oral wound healing. *N Y State Dent J* 1995;61:40-43.
- Damante CA, Gregghi SWL, Sant'Ana AC, Passanezi E, Taga R. Histomorphometric study of the healing of human oral mucosa after gingivoplasty and low-level laser therapy. *Lasers Surg Med* 2004;35:377-384.
- Masse J-F, Landry RG, Rochette C, Dufour L, Morency R, D'Aoust P. Effectiveness of soft laser treatment in periodontal surgery. *Int Dent J* 1993;43:121-127.
- Crespi R, Covani U, Margarone JE, Andreana S. Periodontal tissue regeneration in Beagle dogs after laser therapy. *Lasers Surg Med* 1997;21:395-402.
- Hendrick DA, Meyers A. Wound healing after laser surgery. *Otolaryngol Clin North Am* 1995;28:969-986.
- Zaffe D, Vitale MC, Martignone A, Scarpelli F, Botticelli AR. Morphological, histochemical, and immunocytochemical study of CO₂ and Er:YAG laser effect on oral soft tissues. *Photomed Laser Surg* 2004;22:185-189.
- White JM, Gekelman D, Shin K-B, et al. Lasers and dental soft tissues: Reflections on our years of research. In: Ishikawa I, Frame JW, Aoki A, eds. *Lasers in Dentistry*. Amsterdam: Elsevier Science; 2003:13-19.
- Eriksson RA, Albrektsson T. Temperature threshold levels for heat induced bone tissue injury: A vital microscopic study in the rabbit. *J Prosthet Dent* 1983;50:101-107.
- Fontana CR, Kurachi C, Mendonca CR, Bagnato VS. Temperature variation at soft periodontal and rat bone tissues during a medium-power diode laser exposure. *Photomed Laser Surg* 2004;22:519-522.
- Spencer P, Cobb CM, Wieliczka DM, Glaros AG, Morris PJ. Change in temperature of subjacent bone during soft tissue laser ablation. *J Periodontol* 1998;69:1278-1282.
- McDavid VG, Cobb CM, Rapley JW, Glaros AG, Spencer P. Laser irradiation of bone: III. Long-term healing following treatment by CO₂ and Nd:YAG lasers. *J Periodontol* 2001;72:174-182.
- Friesen LR, Cobb CM, Rapley JW, Forgas-Brockman L, Spencer P. Laser irradiation of bone: II. Healing response following treatment by CO₂ and Nd:YAG lasers. *J Periodontol* 1999;70:75-83.
- Sasaki KM, Aoki A, Ichinose S, Ishikawa I. Ultrastructural analysis of bone tissue irradiated by Er:YAG laser. *Lasers Surg Med* 2002;31:322-332.
- Sasaki KM, Aoki A, Ichinose S, Yoshino T, Yamada S, Ishikawa I. Scanning electron microscopy and Fourier transformed infrared spectroscopy analysis of bone removal using Er:YAG and CO₂ lasers. *J Periodontol* 2002;73:643-652.
- Kimura Y, Yu DG, Fujita A, Yamashita A, Murakami Y, Matsumoto K. Effects of erbium, chromium:YSGG laser irradiation on canine mandibular bone. *J Periodontol* 2001;72:1178-1182.
- Wang X, Zhang C, Matsumoto K. In vivo study of the healing processes that occur in the jaws of rabbits following perforation by an Er, Cr:YSGG laser. *Lasers Med Sci* 2005;20:21-27.
- Aoki A, Sasaki KM, Watanabe H, Ishikawa I. Lasers in nonsurgical periodontal therapy. *Periodontol* 2000;2004;36:59-97.
- Spencer P, Cobb CM, McCollum MH, Wieliczka DM. The effects of CO₂ laser and Nd:YAG with and without water/air surface cooling on tooth root structure: Correlation between FTIR spectroscopy and histology. *J Periodontol Res* 1996;31:453-462.
- Pant V, Dixit J, Agrawal AK, Seth PK, Pant AB. Behavior of human periodontal ligament cells on CO₂ laser irradiated dental root surfaces: An *in vitro* study. *J Periodontol Res* 2004;39:373-379.
- Crespi R, Barone A, Covani U, Ciaglia RN, Romanos GE. Effects of CO₂ laser treatment on fibroblast attachment to root surfaces. A scanning electron microscopy analysis. *J Periodontol* 2002;73:1308-1312.

40. Fayad MI, Hawkinson R, Daniel J, Hao J. The effect of CO₂ laser irradiation on PDL cell attachment to resected root surfaces. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004;97:518-523.
41. Gopin BW, Cobb CM, Rapley JW, Killoy WJ. Histologic evaluation of soft tissue attachment to CO₂ laser-treated root surfaces: An in vivo study. *Int J Periodontics Restorative Dent* 1997;17:316-325.
42. Barone A, Covani U, Crespi R, Romanos GE. Root surface morphological changes after focused versus defocused CO₂ laser irradiation: A scanning electron microscopy analysis. *J Periodontol* 2002;73:370-373.
43. Misra V, Mehrotra KK, Dixit J, Maitra SC. Effect of a carbon dioxide laser on periodontally involved root surfaces. *J Periodontol* 1999;70:1046-1052.
44. Israel M, Cobb CM, Rossmann JA, Spencer P. The effects of CO₂, Nd:YAG and Er:YAG lasers with and without surface coolant on tooth root surfaces. An in vitro study. *J Clin Periodontol* 1997;24:595-602.
45. Trylovich DJ, Cobb CM, Pippin DJ, Spencer P, Killoy WJ. The effects of the Nd:YAG laser on in vitro fibroblast attachment to endotoxin-treated root surfaces. *J Periodontol* 1992;63:626-632.
46. Spencer P, Trylovich DJ, Cobb CM. Chemical characterization of lased root surfaces using Fourier transform infrared photoacoustic spectroscopy. *J Periodontol* 1992;63:633-636.
47. Thomas D, Rapley J, Cobb C, Spencer P, Killoy W. Effects of the Nd:YAG laser and combined treatments on in vitro fibroblast attachment to root surfaces. *J Clin Periodontol* 1994;21:38-44.
48. Gašpirc B, Skalerič U. Morphology, chemical structure and diffusion processes of root surface after Er:YAG and Nd:YAG laser irradiation. *J Clin Periodontol* 2001;28:508-516.
49. Morlock BJ, Pippin DJ, Cobb CM, Killoy WJ, Rapley JW. The effect of Nd:YAG laser exposure on root surfaces when used as an adjunct to root planing: An in vitro study. *J Periodontol* 1992;63:637-641.
50. Chen Y-J, Jeng J-H, Yao C-CJ, Chen M-H, Hou L-T, Lan W-H. Long-term effect of pulsed Nd:YAG laser irradiation on cultured human periodontal fibroblasts. *Lasers Surg Med* 2005;36:225-233.
51. Ito K, Nishikata J, Murai S. Effects of Nd:YAG laser radiation on removal of a root surface smear layer after root planing: A scanning electron microscopic study. *J Periodontol* 1993;64:547-552.
52. Wilder-Smith P, Arrastia AM, Schell MJ, Liaw LH, Grill G, Berns MW. Effect of Nd:YAG laser irradiation and root planing on the root surface: Structural and thermal effects. *J Periodontol* 1995;66:1032-1039.
53. Sakurai Y, Yamaguchi M, Abiko Y. Inhibitory effect of low-level laser irradiation of LPS-stimulated prostaglandin E₂ production and cyclooxygenase-2 in human gingival fibroblasts. *Eur J Oral Sci* 2000;108:29-34.
54. Pourzarandian A, Watanabe H, Ruwanpura SMPM, Aoki A, Noguchi K, Ishikawa I. Er:YAG laser irradiation increases prostaglandin E₂ production via the induction of cyclooxygenase-2 mRNA in human gingival fibroblasts. *J Periodontal Res* 2005;40:182-186.
55. Kreisler M, Meyer C, Stender E, Daubländer M, Willershausen-Zönnchen B, d'Hoedt B. Effect of diode laser irradiation on the attachment rate of periodontal ligament cells: An in vitro study. *J Periodontol* 2001;72:1312-1317.
56. Marques MM, Pereira AN, Fujihara NA, Nogueira FN, Eduardo CP. Effect of low-power laser irradiation on protein synthesis and ultrastructure of human gingival fibroblasts. *Lasers Surg Med* 2004;34:260-265.
57. Kreisler M, Christoffers AB, Willershausen B, d'Hoedt B. Effect of low-level GaAlAs laser irradiation on the proliferation rate of human periodontal ligament fibroblasts: An in vitro study. *J Clin Periodontol* 2003;30:353-358.
58. Kreisler M, Al Haj H, Daubländer M, et al. Effect of diode laser irradiation on root surfaces in vitro. *J Clin Laser Med Surg* 2002;20:63-69.
59. Schwarz F, Aoki A, Sculean A, Georg T, Scherbaum W, Becker J. In vivo effects of an Er:YAG laser, and ultrasonic system and scaling and root planing on the biocompatibility of periodontally diseased root surfaces in cultures of human PDL fibroblasts. *Lasers Surg Med* 2003;33:140-147.
60. Schwarz F, Sculean A, Berakdar M, Szathmari L, Georg T, Becker J. In vivo and in vitro effects of an Er:YAG laser, a GaAlAs diode laser, and scaling and root planing on periodontally diseased root surfaces: A comparative histologic study. *Lasers Surg Med* 2003;32:359-366.
61. Frentzen M, Braun A, Aniol D. Er:YAG laser scaling of diseased root surfaces. *J Periodontol* 2002;73:524-530.
62. Crespi R, Barone A, Covani U. Effect of Er:YAG laser on diseased root surfaces: An in vivo study. *J Periodontol* 2005;76:1386-1390.
63. Schoop U, Moritz A, Maleschitz P, et al. The impact of Er:YAG laser irradiation on root surfaces: An in vitro evaluation. *J Oral Laser Appl* 2001;1:35-41.
64. Schwarz F, Pütz N, Georg T, Reich E. Effect of an Er:YAG laser on periodontally involved root surfaces: An in vivo and in vitro SEM comparison. *Lasers Surg Med* 2001;29:328-335.
65. Yamaguchi H, Kobayashi K, Osada R, et al. Effects of irradiation of an Erbium:YAG laser on root surfaces. *J Periodontol* 1997;68:1151-1155.
66. Theodoro LH, Haypek P, Bachmann L, et al. Effect of Er:YAG and diode laser irradiation on the root surface: Morphological and thermal analysis. *J Periodontol* 2003;74:838-843.
67. Feist IS, De Micheli G, Carneiro SRS, Eduardo CP, Miyagi SPH, Marques MM. Adhesion and growth of cultured human gingival fibroblasts on periodontally involved root surfaces treated by Er:YAG laser. *J Periodontol* 2003;74:1368-1375.
68. Schoop U, Moritz A, Kluger W, et al. Changes in root surface morphology and fibroblast adherence after Er:YAG laser irradiation. *J Oral Laser Appl* 2002;2:83-93.
69. Sasaki KM, Aoki A, Ichinose S, Ishikawa I. Morphological analysis of cementum and root dentin after Er:YAG laser irradiation. *Lasers Surg Med* 2002;31:79-85.
70. Folwaczny M, Benner KU, Plasskamp B, Mehl A, Hickel R. Effects of 2.94 micron Er:YAG laser radiation on root surfaces treated in situ: A histological study. *J Periodontol* 2003;74:360-365.
71. Crespi R, Romanos GE, Barone A, Sculean A, Covani U. Er:YAG laser in defocused mode for scaling of periodontally involved root surfaces: An in vitro pilot study. *J Periodontol* 2005;76:686-690.
72. Shafik SS, Kheir AO, Kany F, Omran M. Effect of Nd:YAP laser on dental cementum: A scanning electron microscopic study. *J Oral Laser Appl* 2002;2:95-99.

73. van As G. Erbium lasers in dentistry. *Dent Clin North Am* 2004;48:1017-1059.
74. Coluzzi DJ, Goldstein AJ. Lasers in dentistry: An overview. *Dent Today* 2004;23:120-127.
75. Parker S. The use of lasers in fixed prosthodontics. *Dent Clin North Am* 2004;48:971-998.
76. Flax H. Maximizing esthetic transformations using a closed flap Er, Cr:YSGG modality. *Compend Contin Educ Dent* 2005;26: 172, 174, 176 passim.
77. Flax H. Closed-flap laser-assisted esthetic dentistry using Er:YSGG technology. *Compend Contin Educ Dent* 2004;25: 622, 626, 628-630 passim.
78. Adams TC, Pang PK. Lasers in aesthetic dentistry. *Dent Clin North Am* 2004;48:833-860.
79. Cobb CM, McCawley TK, Killoy WJ. A preliminary in vivo study on the effects of the Nd:YAG laser on root surfaces and subgingival microflora. *J Periodontol* 1992;63:701-707.
80. Ben Hatit Y, Blum R, Severin C, Maquin M, Jabro MH. The effects of a pulsed Nd:YAG laser on subgingival bacterial flora and on cementum: An in vivo study. *J Clin Laser Med Surg* 1996;14:137-143.
81. Gutknecht N, Radufi P, Franzen R, Lampert F. Reduction of specific microorganisms in periodontal pockets with the aid of an Nd:YAG laser – An in vivo study. *J Oral Laser Appl* 2002;2:175-180.
82. Radvar M, Creanor SL, Gilmour WH, et al. An evaluation of the effects of an Nd:YAG laser on subgingival calculus, dentine and cementum. An in vitro study. *J Clin Periodontol* 1995;22:71-77.
83. Meral G, Tasar F, Kocagöz S, Sener C. Factors affecting the antibacterial effects of Nd:YAG laser in vivo. *Lasers Surg Med* 2003;32:197-202.
84. Harris DM, Yessik M. Therapeutic ratio quantifies laser antiseptics: Ablation of *Porphyromonas gingivalis* with dental lasers. *Lasers Surg Med* 2004;35:206-213.
85. Moritz A, Gutknecht N, Doertbudak O, et al. Bacterial reduction in periodontal pockets through irradiation with a diode laser: A pilot study. *J Clin Laser Med Surg* 1997;15:33-37.
86. Nussbaum EL, Lilje L, Mazzulli T. Effects of 810 nm laser irradiation on in vitro growth of bacteria: Comparison of continuous wave and frequency modulated light. *Lasers Surg Med* 2002;31:343-351.
87. Coffelt DW, Cobb CM, Rapley JW, Killoy WJ. Determination of energy density threshold for laser ablation of bacteria: An in vitro study. *J Clin Periodontol* 1997; 24:1-7.
88. Crespi R, Barone A, Covani U. Histologic evaluation of three methods of periodontal root surface treatment in humans. *J Periodontol* 2005;76:476-481.
89. Tucker D, Cobb CM, Rapley JW, Killoy WJ. Morphologic changes following in vitro CO₂ laser treatment of calculus-laden root surfaces. *Lasers Surg Med* 1996; 18:150-156.
90. Ando Y, Watanabe H, Ishikawa I. Bactericidal effect of erbium YAG laser on periodontopathic bacteria. *Lasers Surg Med* 1996;19:190-200.
91. Aoki A, Ando Y, Watanabe H, Ishikawa I. In vitro studies on laser scaling of subgingival calculus with an erbium:YAG laser. *J Periodontol* 1994;65:1097-1106.
92. Folwaczny M, Mehl A, Haffner C, Benz C, Hickel R. Root substance removal with Er:YAG laser radiation at different parameters using a new delivery system. *J Periodontol* 2000;71:147-155.
93. Aoki A, Miura M, Akiyama F, et al. In vitro evaluation of Er:YAG laser scaling of subgingival calculus in comparison with ultrasonic scaling. *J Periodontol Res* 2000;35:266-277.
94. Folwaczny M, Mehl A, Aggstaller H, Hickel R. Antimicrobial effects of 2.94 μm Er:YAG laser radiation on root surfaces: An in vitro study. *J Clin Periodontol* 2002;29:73-78.
95. Eberhard J, Ehlers H, Falk W, Acil Y, Albers HK, Jepsen S. Efficacy of subgingival calculus removal with Er:YAG laser compared to mechanical debridement: An in situ study. *J Clin Periodontol* 2003;30: 511-518.
96. Waerhaug J. Healing of the dento-epithelial junction following subgingival plaque control. II. As observed on extracted teeth. *J Periodontol* 1978;49:119-134.
97. Caton JG, Zander HA. The attachment between tooth and gingival tissues after periodic root planing and soft tissue curettage. *J Periodontol* 1979;50:462-466.
98. Beaumont RH, O'Leary TJ, Kafrawy AH. Relative resistance of long junctional epithelial adhesions and connective tissue attachments to plaque-induced inflammation. *J Periodontol* 1984;55:213-223.
99. Magnusson I, Runstad L, Nyman S, Lindhe J. A long junctional epithelium – A locus minoris resistentiae in plaque infection? *J Clin Periodontol* 1983;10:333-340.
100. Borrajo JL, Varela LG, Castro GL, Rodriguez-Nunez I, Torreira MG. Diode laser (980 nm) as adjunct to scaling and root planing. *Photomed Laser Surg* 2004;22:509-512.
101. Neill ME, Mellonig JT. Clinical effects of the Nd:YAG laser for combination periodontitis therapy. *Pract Periodontics Aesthet Dent* 1997;9(Suppl. 6):1-5.
102. Miyazaki A, Yamaguchi T, Nishikata J, et al. Effects of Nd:YAG and CO₂ laser treatment and ultrasonic scaling on periodontal pockets of chronic periodontitis patients. *J Periodontol* 2003;74:175-180.
103. Schwarz F, Sculean A, Georg T, Reich E. Periodontal treatment with an Er:YAG laser compared to scaling and root planing. A controlled clinical study. *J Periodontol* 2001;72:361-367.
104. Schwarz F, Sculean A, Berakdar M, Georg T, Reich E, Becker J. Clinical evaluation of an Er:YAG laser combined with scaling and root planing for non-surgical periodontal treatment. A controlled, prospective clinical study. *J Clin Periodontol* 2003; 30:26-34.
105. Schwarz F, Sculean A, Berakdar M, Georg T, Reich E, Becker J. Periodontal treatment with an Er:YAG laser or scaling and root planing. A 2-year follow-up split-mouth study. *J Periodontol* 2003;74:590-596.
106. Sculean A, Schwarz F, Berakdar M, Romanos GE, Arweiler NB, Becker J. Periodontal treatment with an Er:YAG laser compared to ultrasonic instrumentation: A pilot study. *J Periodontol* 2004;75: 966-973.
107. Schwarz F, Sculean A, Georg T, Becker J. Clinical evaluation of the Er:YAG laser in combination with an enamel matrix protein derivative for the treatment of intrabony periodontal defects: A pilot study. *J Clin Periodontol* 2003;30:975-981.
108. Sculean A, Schwarz F, Berakdar M, Windisch P, Arweiler NB, Romanos GE. Healing of intrabony defects following surgical treatment with or without an Er:YAG laser. *J Clin Periodontol* 2004;31:604-608.
109. El Yazami H, Azehoui N, Ahariz M, Rey G, Sauvete E. Periodontal evaluation of an Nd:YAP laser combined with scaling and root planing for nonsurgical

- periodontal treatment. A clinical evaluation. *J Oral Laser Appl* 2004;4:97-102.
110. Ambrosini P, Miller N, Briancon S, Gallina S, Penaud J. Clinical and microbiological evaluation of the effectiveness of the Nd:YAP laser for the initial treatment of adult periodontitis: A randomized controlled study. *J Clin Periodontol* 2005;32:670-676.
111. Noguchi T, Sanaoka A, Fukuda M, Suzuki S, Aoki T. Combined effects of Nd:YAG laser irradiation with local antibiotic applications into periodontal pockets. *J Int Acad Periodontol* 2005;7:8-15.
112. Rydén H, Persson L, Preber H, Bergström J. Effect of low level energy laser irradiation on gingival inflammation. *Swed Dent J* 1994;18:35-41.
113. Yilmaz S, Kuru B, Kuru L, Noyan Ü, Argun D, Kadir T. Effect of gallium arsenide diode laser on human periodontal disease: A microbiological and clinical study. *Lasers Surg Med* 2002;30:60-66.
114. Qadri T, Miranda L, Tuner J, Gustafsson A. The short-term effects of low-level lasers as adjunct therapy in the treatment of periodontal inflammation. *J Clin Periodontol* 2005;32:714-719.
115. Moritz A, Schoop U, Goharkhay K, et al. Treatment of periodontal pockets with a diode laser. *Lasers Surg Med* 1998;22:302-311.
116. Finkbeiner RL. The results of 1328 periodontal pockets treated with the argon laser: Selective pocket thermolysis. *J Clin Laser Med Surg* 1995;13:273-281.
117. Radvar M, MacFarlane TW, MacKenzie D, Whitters CJ, Payne AP, Kinane DF. An evaluation of the Nd:YAG laser in periodontal pocket therapy. *Br Dent J* 1996;180:57-62.
118. Liu C-M, Hou L-T, Wong M-Y, Lan W-H. Comparison of Nd:YAG laser versus scaling and root planing in periodontal therapy. *J Periodontol* 1999;70:1276-1282.
119. Sjöstrom L, Friskopp J. Laser treatment as an adjunct to debridement of periodontal pockets. *Swed Dent J* 2002;26:51-57.
120. Harris DM, Gregg RH II, McCarthy DK, Colby LE, Tilt LV. Laser-assisted new attachment procedure in private practice. *Gen Dent* 2004;52:396-403.

Correspondence: Dr. Charles M. Cobb, Department of Periodontics, School of Dentistry, University of Missouri, 424 W. 67th Terrace, Kansas City, MO 64113. Fax: 816/444-8673; e-mail: cobbcc@umkc.edu.

Accepted for publication January 16, 2006.